



REDUCING HAZARDOUS DRINKING AMONG WOMEN WHO HAVE SERVED IN THE UNITED KINGDOM ARMED FORCES: A RANDOMISED CONTROLLED TRIAL OF THE DRINKSRATION APP

CIMVHR FORUM 2026 | HALIFAX, NOVA SCOTIA

Dr Dan Leightley

King's College London (UK)

daniel.leightley@kcl.ac.uk



PRESENTER

- Daniel Leightley is employed by King's College London.
- He is a member of the UK Armed Forces Reserve.
- He was involved in the development and evaluation of the DrinksRation intervention.
- No relevant commercial or financial conflicts of interest to declare.



PRESENTER

- No commercial support was received for this study or presentation.
- This study was funded by the UK Office for Veterans' Affairs (grant G2-SCH-2022-11-12241).
- Earlier development of DrinksRation was supported by the UK Medical Research Council and Forces in Mind Trust.
- Combat Stress has provided support towards computing costs associated with DrinksRation.



PRESENTER

- The presenter's involvement in the development of DrinksRation represents a potential source of bias.
- The study used a randomised, participant-blinded design with pre-specified outcomes and analyses.
- Results reflect evaluation of the intervention as a whole, rather than the effectiveness of individual intervention components.



By the end of the presentation, participants should be able to:

- | | | |
|-----------|------------------|---|
| 01 | Describe | why alcohol support should address the experiences of women who have served. |
| 02 | Interpret | the trial's 84-day alcohol and AUDIT outcomes, including uncertainty and attrition. |
| 03 | Discuss | implications for military and veteran services, and the limits of transferability. |



A health need, barriers to support, and an evidence gap...

Health need

Women who have served report higher levels of hazardous drinking than women in the general population.

Access to support

Stigma and barriers to help-seeking can limit access to appropriate support.

Evidence gap

Earlier DrinksRation research did not address women's experiences and needs.

RESEARCH QUESTION

Does DrinksRation reduce weekly alcohol consumption more than standard digital advice over 84 days?





DrinksRation

Save Money


By drinking less with DrinksRation.



Download DrinksRation on iOS


Goals

Set Goals to help you cut down.



Log

Record any time.




Insights

Insights about your drinking.





The core was retained; content and presentation were co-designed.



Participants involved in co-design or pre-trial testing were excluded from the trial.



Two-arm, participant-blinded, 1:1 randomised trial; fully remote

ELIGIBILITY

Women aged 18+; UK resident

At least one paid day of UK Armed Forces service

An iOS or Android smartphone

Consumed more than 14 UK units per week

OUTCOMES

PRIMARY

7-day Timeline Follow-Back:
weekly alcohol units

SECONDARY

Alcohol Use Disorders

Identification Test (AUDIT) score

Baseline and days 28, 56 and 84 | Primary endpoint: day 84 (12 weeks)



An interactive DrinksRation versus web-based alcohol advice

DRINKSRATION

Smartphone app

Diary, feedback and goal setting

Content for women who have served

Personalised messages and prompts

Questionnaires within the app



Download DrinksRation on iOS

BEALCOHOLSMART

Web app

NHS advice and 7-day unit calculator

Emails at enrolment and assessments


Questionnaires through Qualtrics

Less interactive content and contact



BEALCOHOLSMART

PARTICIPANT FLOW AND ANALYSED SAMPLE



Substantial loss occurred before inclusion in the analysis.

47.4 years

Mean age

27.2 units


Baseline weekly use

PROBABLE MENTAL HEALTH CONDITIONS

Depression: 52% | Anxiety: 39% | PTSD: 31%



Greater reduction in reported weekly alcohol consumption at day 84



ADJUSTED DAY-84 DIFFERENCE

-11.6

UK units per week


95% CI: -19.7 to -3.6

$p=0.005$

About 6.9 fewer Canadian
standard drinks per week



AUDIT scores were also lower in the DrinksRation group at day 84.



ADJUSTED DAY-84 DIFFERENCE

−3.9

AUDIT points

95% CI: −6.9 to −1.0

$p=0.010$



More recorded engagement and higher usability ratings for the app

MEDIAN WEEKS ACTIVE

11 versus 1

DrinksRation versus BeAlcoholSmart

MEAN USABILITY AT DAY 28

6.22 versus 5.22

Out of 7; app $n=39$, control $n=30$

No serious safety concerns or technical issues were reported.



Results favour DrinksRation, but...

Selection and missing data

88 women contributed outcome data. The model assumes outcomes are missing at random.

Outcomes and duration

Self-reported alcohol use. Effects beyond 84 days and on clinical harm are not established.


Unequal intervention packages

Content, functionality and contact differed. Tailoring could not be isolated.

CONCLUSIONS



- DrinksRation significantly reduced weekly alcohol consumption and AUDIT scores at 84 days.
- Engagement and usability were high, with no safety or technical concerns.
- Findings support digital alcohol interventions for women who have served in the Armed Forces.
- Effects reflect the whole DrinksRation package, and substantial attrition and short follow-up mean further evaluation is needed.



THANK YOU



Research Team

Daniel Leightley
Grace Williamson
Ewan Carr
Rhea Varghese
Simon Dymond
Kate King
Amos Simms
Dominic Murphy
Laura Goodwin
Nicola T. Fear

Reducing alcohol consumption in UK Armed Forces veterans: feasibility of using personalized push notifications with AI

Daniel Leightley, Charlotte Williamson, Iain J. Marshall, Vasa Curcin, Roberto J. Rona, Dominic Murphy, Nicola T. Fear and Laura Goodwin. *PLOS Digital Health*, 2026.



In memory of Professor Nicola T. Fear, with gratitude for her exceptional contribution to this work and to military health research.



Rhead R, MacManus D, Jones M, Greenberg N, Fear NT, Goodwin L. Mental health disorders and alcohol misuse among UK military veterans and the general population: a comparison study. *Psychological Medicine*. 2020;1-11. doi:10.1017/S0033291720001944.

Godier-McBard LR, Cable G, Wood AD, Fossey M. Gender differences in barriers to mental healthcare for UK military veterans: a preliminary investigation. *BMJ Military Health*. 2022;168(1):70-75. doi:10.1136/bmjmilitary-2020-001754.

Leightley D, Williamson C, Rona RJ, Carr E, Shearer J, Davis JP, et al. Evaluating the efficacy of the Drinks:Ration mobile app to reduce alcohol consumption in a help-seeking military veteran population: randomized controlled trial. *JMIR mHealth and uHealth*. 2022;10(6):e38991. doi:10.2196/38991.

Williamson G, Carr E, Fear NT, Dymond S, King K, Simms A, et al. Digital therapeutic intervention for women in the UK Armed Forces who consume alcohol at a hazardous or harmful level: protocol for a randomized controlled trial. *JMIR Research Protocols*. 2023;12:e51531. doi:10.2196/51531.

Zhou L, Bao J, Setiawan IMA, Saptono A, Parmanto B. The mHealth App Usability Questionnaire (MAUQ): development and validation study. *JMIR mHealth and uHealth*. 2019;7(4):e11500. doi:10.2196/11500.